



0 184 692
A1

EUROPEAN PATENT APPLICATION

(51) Int. Cl.: **C 07 F 7/18**, C 07 F 7/04,
C 07 F 7/02, C 07 F 7/21

② Date of filing: 19.11.85

⑦ Applicant: **UNION CARBIDE CORPORATION**, 39 Old Ridgebury Road, Danbury Connecticut 06817 (US)


(72) Inventor: Quirk, Jennifer Maryann, KB-208 Harris Road, Bedford Hills 10507 (US)
Inventor: Kozak, Linda Kathy, 11 Madison Avenue, Ossining 10562 (US)
Inventor: Kanner, Bernard, 50 Robin Lane, West Nyack 10994 (US)

**74 Representative: Wuesthoff, Franz, Dr.-Ing. et al,
Patentanwlte Wuesthoff -v. Pechmann-Behrens-Goetz
Schweigerstrasse 2, D-8000 Mnchen 90 (DE)**


57) O-silylated ketene acetals and enol ethers are prepared via the 1,4-hydrosilylation of a hydrosilylation composition selected from the group consisting of aminosilanes, siloxanes and alkylalkoxysilanes with an α,β -unsaturated carbonyl in the presence of a rhodium catalyst.

These compounds consist essentially of:


(a) O-aminosilylated ketene acetals and enol ethers of the general formula:



(b) O-siloxanesilylated ketene acetals and enol ethers of the general formulae:



(c) O-alkoxysilylated ketene acetals and enol ethers of the general formula:



wherein R, R¹ and R² are individually alkyl or aryl group and R can be hydrogen; R³, R⁴, R⁵, R⁶, R⁷, and R¹⁰ are individually alkyl groups; R⁸ is hydrogen or an alkyl or aryl, Z is selected from hydrogen, an alkyl group, an aryl group, an alkoxy group and an aryloxy group

and a is 0 or 1, b 0 or 1, y has a value from 1 to 500, z has a value from 0 to 499, and, the sum of $y+z$ equals from 1 to 500 with the proviso that if z is 0, $a + b$ must equal at least 1; x is 1, 2 or 3; and v is 0, 1 or 2.

PATENTANWÄLTE
WUESTHOFF-v. PECHMANN-BEHRENS-GOETZ
EUROPEAN PATENT ATTORNEYS

Union Carbide Corp.
Our File: EP-59 902

0184692
DR.-ING. FRANZ WUESTHOFF
DR. PHIL. FREDA WUESTHOFF (1927-1956)
DIPLOM.-ING. GERHARD PULS (1952-1971)
DIPLOM.-CHEM. DR. E. FREIHERR VON PECHMANN
DR.-ING. DIETER BEHRENS
DIPLOM.-ING.; DIPLOM.-WIRTSCH.-ING. RUPERT GOETZ
D-8000 MÜNCHEN 90
SCHWEIGERSTRASSE 2
TELEFON: (089) 66 20 51
TELEGRAMM: PROTECTPATENT
TELEX: 5 24 070
TELEFAX: VIA (089) 2 71 60 63 (111)

O-SILYLATED KETENE ACETALS AND ENOL ETHERS
AND METHOD OF PREPARATION

BACKGROUND OF THE INVENTION

The present invention generally relates to
O-silylated ketene acetals and enol
ethers and a process for preparing the same. More
specifically, the present invention is concerned
with those O-silylated ketene acetals and enol
ethers which are prepared via the 1,4 hydrosilation
of a hydrosilation composition selected from
of aminosilanes, siloxanes and
alkylalkoxysilanes with an α,β -unsaturated
carbonyl compound.

THE PRIOR ART

In U.S.-A., 4,417,034 and 4,414,372
it was reported that a variety of O-silylated ketene
acetals have utility as initiators for group
transfer polymerization. O-silylated ketene acetals
and enol ethers can be prepared via several known
synthetic routes, among which include hydrosilation.

The hydrosilation reaction was discovered
in 1947 and over the years has become one of the
best known and most widely practiced reactions in
organosilicon chemistry, including its use in a wide
variety of large scale commercial applications. It
has also been the subject of several extensive

reviews, see for instance: Organic Insertion Reactions of Group II Elements, Consultants Bureau, N. Y., 1966; Organometallic Compounds of the Group IV Elements, Dekker, N. Y., 1968, Vol. I; Preparation of Carbofunctional Organosilanes by an Addition Reaction, Moscow, 1971; Russ. Chem Rev., 46, 264 (1977); and J. Organometal Chem. Library 5, 1977, pg. 1-179.

In operating a hydrosilation reaction, various transition metals are known to be effective catalysts. U.S. - A - . 2,823,218 teaches chloroplatinic acid, a soluble form of platinum, as a particularly effective hydrosilation catalyst.

The prior art involving hydrosilations by aminosilanes is concentrated, in large measure, in a series of papers published between 1968-1970. Russian workers reported that disilazanes such as $\text{HN}(\text{SiMe}_2\text{H})_2$ and cyclosilazanes such as $(\text{HNMe}_2\text{H})_4$ underwent slow hydrosilation: Izv. Akad. Nauk SSSR, Ser. Khim. 1968, pg 351-6; and 1969, pg 1539-45; Zhur. Obshch. Khim, 38, 655, (1968); and 40, 1082, (1970). These reports were confirmed by workers at Dow Corning who went on to demonstrate that mono-, bis-, and tris- (dialkylamino) silanes showed no reactivity for hydrosilation J. Org. Chem. 35, 3879 (1970).

There appeared to be little hydrosilation potential for aminosilanes until a Russian paper disclosed reactions of hydrosilazanes with arylacetylenes, Soobshch Akad. Nauk Gruz. SSR, 84, 381 (1976), and, Tezisy Dokl-Vses. Konf. Khim. Atsetilena, 5th, 1975, 172.

Similarly, it was reported in the literature that triethylsilane undergoes a 1,4-hydrosilation with methacrylates using $\text{RhCl}(\text{PPH}_3)_3$ as a catalyst, Chem. Pharm. Bull. Japan, 22 (1974) 2767, and, J. Organomet. Chem 111 (1976) 43. However, following these teachings a mixture of 1,4-and 1,2-hydrosilated products are formed when acrylates are employed in the reaction.

Despite this background of art there continued to exist a need for a one-step preparation of O-silylated ketene acetals and enol ethers which would primarily yield the 1,4-hydrosilated product. This need was significantly increased in view of the discovery that such materials may find utility as initiators in group transfer polymerization. The primary object of this invention was to provide a route to O-silylated ketene acetals and enol ethers as would yield the 1,4-hydrosilated product. Another object of the present invention was to arrive at new classes of O-silylated ketene acetals and enol ethers which may be useful as initiators in group transfer polymerization.

SUMMARY OF THE INVENTION


The present invention provides

O-silylated ketene acetals and enol ethers and a one-step method of preparing such O-silylated ketene acetals and enol ethers which primarily yields the 1,4-hydrosilated product. The O-silylated ketene acetals and enol ethers are those prepared via a 1,4-hydrosilation of a hydrosilation

composition selected from aminosilanes, siloxanes, and alkylalkoxysilanes with an α,β -unsaturated carbonyl compound in the presence of a rhodium catalyst.

DETAILED DESCRIPTION OF THE INVENTION

Aminosilanes useful in the hydrosilation include those represented by the general formula:




wherein R, R¹ and R² are individually alkyl or aryl groups containing from 1 to 8 carbon atoms R may also be hydrogen and wherein x has a value of 1, 2 or 3. These aminosilanes may be purchased commercially or prepared from any of a variety of known techniques, such as described in Eaborn, Organosilicon Compounds, Academic Press Inc., New York, 1960, p. 339.

Suitable aminosilanes include,
 . dimethylaminodimethylsilane,
bis(dimethylamino)methylsilane, tris(dimethylamino)
silane, dimethylaminomethylethylsilane,
diethylaminomethylpropylsilane,
methylaminomethylethylsilane,

ethylaminomethylethylsilane,
 phenylaminomethylethylsilane;
 diphenylaminomethylphenylsilane,
 dibenzylaminoethylphenylsilane,
 diphenylaminodimethylsilane,
 bis(diethylamino)ethylsilane,
 bis(methylamino)ethylsilane,
 bis(benzylamino)methylsilane,
 tris(diethylamino)silane, tris(piperidino)silane,
 tris(cyclohexylamino)silane. Preferably the
 aminosilane is such that R, R¹ and R² are either
 methyl, ethyl or phenyl and x has a value of 3.
 Preferably the silane is tris(dimethylamino)silane.

Siloxanes useful in the hydrosilation
 include those represented by the general formula:



where in R³, R⁴, R⁵ and R⁶ are individually
 alkyl groups containing from 1 to 8 carbon atoms,
 and a has a value of 0 or 1, b has a value of 0 or
 1, y has a value from 1 to 500, z has a value from 0
 to 499, and, the sum of y+z equals from 1 to 500
 with the proviso that if z is 0, a+b must equal at
 least one. The siloxanes may be purchased
 commercially or prepared from any of a variety of
 known techniques, such as described in Eaborn,
Organosilicon Compounds, Academic Press Inc, New
 York, 1960, p. 228.

Suitable siloxanes include,

, bis(trimethylsilyloxy)methylsilane
 1,1,3,3-tetramethyldisiloxane, and

heptamethylcyclotetrasiloxane. Most other siloxanes are mixtures of all types of siloxanes. Preferably the siloxane is such that R^3 , R^4 , R^5 , R^6 are all methyl groups.


Alkylalkoxysilanes useful in the hydrosilation include those represented by the general formula set forth below:



wherein R^7 and R^{10} may be the same or different throughout the silane and are individually an alkyl group containing from 1 to 8 carbon atoms and v is 0, or 1 or 2. The alkylalkoxysilanes may be purchased commercially or prepared by a variety of known techniques such as are described for example in U.S. - A - 4,395,564. It is preferred that the alkylalkoxysilanes be prepared from secondary or tertiary alcohols and v is 0.

Suitable alkylalkoxysilanes include,
 , triisopropoxysilane,
 trisecbutoxysilane, tritertbutoxysilane,
 triethoxysilane, trimethoxysilane,
 methyldiethoxysilane, dimethylethoxysilane,
 methyldiisopropoxysilane, dimethyltertbutoxysilane,
 methyldisecbutoxysilane, and dimethylmethoxysilane.

The α,β -unsaturated monomers the hydrosilation is run with is represented by the general formula:



(IV)

wherein R^8 can be hydrogen or an alkyl or aryl group having from 1 to 8 carbon atoms and Z is selected from hydrogen, an alkyl group, an aryl group, an alkoxy group and an aryloxy group containing from 1 to 15 carbon atoms, preferably from 1 to 8 carbon atoms. The α, β -unsaturated carbonyl compound such as acrylates or methacrylates can be purchased commercially or may be prepared by any of a variety of known techniques.

α, β -unsaturated carbonyls which are suitable for use in the practice of this invention are, in general, known compounds and include,

the following: methyl methacrylate; butyl methacrylate; lauryl methacrylate; ethyl acrylate; butyl acrylate; 2-ethylhexyl methacrylate; 2-(dimethylamino)ethyl methacrylate; 2-(dimethylamino)ethyl acrylate; 3,3-dimethoxypropyl acrylate; 3-methacryloxypropyl acrylate; 2-acetoxyethyl methacrylate; p-tolyl methacrylate; 2,2,3,3,4,4,4-heptafluorobutyl acrylate; ethyl 2-cyanoacrylate; N,N-dimethyl acrylamide; ethyl 2-cyanoacrylate; 4-fluorophenyl acrylate; 2-methylacryloxyethyl acrylate and linoleate; propyl vinyl ketone; methyl vinyl ketone; phenyl vinyl ketone; ethyl vinyl ketone; acrolein, methacrolein; ethyl 2-chloroacrylate; glycidyl methacrylate; 3-methoxypropyl methacrylate;

2-[(1-propenyl)oxy]ethyl methacrylate and acrylate; phenyl acrylate; 2-(trimethylsiloxy)ethyl methacrylate; and allyl acrylate and methacrylate. Preferred monomers include methyl methacrylate; glycidyl methacrylate; ethyl acrylate; butyl acrylate; 2-(trimethylsiloxy)ethyl methacrylate; 2-methacryloxyethyl acrylate; 2-acetoxyethyl methacrylate; and 2-(dimethylamino)ethyl methacrylate. Methyl methacrylate is most preferred.

To obtain primarily (more than 50% and usually more than 75%) the 1,4-hydrosilation product the hydrosilation reaction takes place in the presence of a rhodium catalyst. The rhodium catalyst may be used neat; on a support such as carbon black or alumina; or in a solvent as a soluble compound of rhodium, i.e. as rhodium trichloride, rhodium (III) 2,4-pentanedionate and complexes of rhodium with phosphines, such as tris(triphenylphosphine) rhodium chloride and tris(triphenylphosphine) rhodium carbonyl hydride. The rhodium catalyst is available commercially from Johnson Mathey. The concentration of the catalyst is normally between 0.000010-0.05 mole% with respect to the hydrosilation composition, it is preferred to use no more catalyst than required to obtain the necessary reaction due to the significant costs associated with catalysts of this type. However, impurities common in many α,β -unsaturated carbonyl compounds may necessitate higher concentrations.


The hydrosilation should be run at temperatures greater than 40°C with the optimum

being between 70°C and 150°C. Although the upper temperature limit is not critical, the reaction should be below the decomposition point of the starting materials or products. The hydrosilation reaction may be carried out with or without solvent. In most cases there is no advantage to using a solvent. However, in cases where a solvent is desirable for reasons such as solubility or temperature control, a solvent may be used. Suitable solvents are hydrocarbons such as octane, xylene or triisopropylbenzene.


The order of addition of the reactants is not important, although normally, the rhodium catalyst is added to the hydrosilation composition, then heat is applied, followed by the addition of the α,β -unsaturated carbonyl compound. A stoichiometric ratio of 1 : 1 for the hydrosilation composition to the α,β -unsaturated carbonyl compound is preferred, although it is possible to alter this ratio slightly without adversely affecting the reaction. Any such change will, however, affect the reaction's yield.

The O-silylated ketene acetal or enol ether which the 1,4-hydrosilation produces is represented by the following general formulae:

- A. When the hydrosilation composition is an aminosilane:




B. When the hydrosilation composition is an siloxane:



C. When the hydridosilane composition is an alkylalkoxysilane:

(VII)



The O-silylated ketene acetals and enol ethers find utility as initiators for group transfer polymerization. Group-transfer polymerization is a method for the controlled formation of polymers from α , β -unsaturated carbonyl compounds, i.e. esters, ketones, carboxamides, and nitriles via a sequential conjugate (Michael type) addition process. The technique employs silyl ketene acetals as initiators and is based on the known chemistry of the conjugate Michael addition of these silyl reagents to α , β -unsaturated carbonyl compounds. This reaction has been extensively used in organic synthesis in recent years. The innovation contributed by DuPont scientists was the discovery of catalysts which promote the sequential conjugate addition of immediate silyl ketene acetals to α , β -unsaturated carbonyl compounds, i.e.

esters, ketones, carboxamides, or nitriles to form polymers.

The process is termed group-transfer polymerization because it proceeds via transfer of the silyl group from the initiator and each intermediate ketene acetal to the unsaturated heteroatom (e.g., carbonyl oxygen of an ester) of the next monomer molecule to be added to the polymer chain. Thus, the chain ends of all the intermediate molecules are reactive, i.e., the polymers are "living", and the reaction continues until all the α , β -unsaturated carbonyl is consumed or the polymerization is terminated by addition of a reagent (e.g., a protic solvent) which cleaves the silyl group from the polymer molecule.

Group-transfer polymerization will be useful to control the molecular weight of acrylic polymers and thus may find utility in the production of acrylic automobile finishes and related products.

Whereas the exact scope of the instant invention is set forth in the appended claims, the following specific examples illustrate certain aspects of the present invention and, more particularly, point out methods of evaluating the same.

All parts and percentages are by weight unless otherwise specified.

Example 1

A 100 ml 3-neck flask was equipped with a stirring bar, thermometer, 10 ml addition funnel and

reflux condenser topped with a nitrogen inlet tube. The flask was charged with 15.7g (0.07 mol) bis(trimethylsilyloxy)methylsilane, 22.5g xylene, 1.0g phenothiazine and 200 ppm (23.64 mg) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to reflux, at which time 7.1g (0.07 mol) methyl methacrylate was added dropwise to the reaction over a period of 1-2 hours. The reaction was then cooled to room temperature and GC, GC/MS and NMR analyses confirmed the structure to be 1,1-dimethyl-2-methoxy-2-bis(trimethylsilyloxy)methylsiloxylethylene. The product was distilled at 58°C/53 mbar (40 mm Hg).

Example 2

The reaction was run as described in Example 1 with the exception that methyl acrylate was used instead of methyl methacrylate. GC analysis showed that one major product had been formed. GC/MS and NMR confirmed the structure to be 1-methyl-2-methoxy-2-bis(trimethylsilyloxy)methylsiloxylethylene.

Example 3

The reaction was run as described in Example 1 with the exception that the starting hydrosiloxane used was heptamethylcyclotetrasiloxane instead of bis(trimethylsilyloxy)methylsilane. A single major product was formed in approximately 80% yield by GC analysis. GC/MS and NMR confirmed that this product was 1,1-dimethyl-2-methoxy-2-heptamethylcyclotetrasiloxylethylene.

Example 4

A 100 ml 3-neck flask was equipped with a stirring bar, thermometer, 50 ml addition funnel and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 15.0g (0.112 mole), 1,1,3,3-tetramethyldisiloxane, 37.4g xylene, 1.5g phenothiazine and 200 ppm (22.2mg) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to reflux at which time 22.4g (0.224 mol) methyl methacrylate was added dropwise to the reaction over a period of 2-3 hours. The reaction was then cooled to room temperature and GC analysis showed that one major product was formed in greater than 90% yield. GC/MS and NMR confirmed that the structure of the product was 1,3-bis(1,1-dimethyl-2-methoxy-2-ethenoxy)-1,1,3,3-tetramethyldisiloxane.

Example 5

A 250 ml 3-neck flask was equipped with a stirring bar, thermometer, 50 ml addition funnel and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 30.9g (0.15 mol) triisopropoxysilane, 45.9g xylene, 1.82g phenothiazine and 200 ppm (45.78 mg) $\text{RhCl}(\text{PPh}_3)_3$. The reaction was heated to reflux at which time 15.0g (1.5 mol) methyl methacrylate was added dropwise to the reaction mixture over a period of 1-2 hours which was then heated an additional 5 hours. The reaction was then cooled to room temperature and GC analysis showed that one major product has been formed in greater than 80% yield. GC/MS and NMR confirmed the structure to be

1,1-dimethyl-2-methoxy-2-triisopropoxysiloxyethylene. The product was distilled at 110°C/13 mbar (10 mm Hg).

Example 6

The reaction was run as describe in Example 5 with the exception that tritertbutoxysilane was used as the starting silane instead of triisopropoxysilane. GC analysis showed that one major product has been formed in greater than 90% yield. GC/MS and NMR confirmed the structure to be 1,1-dimethyl-2-methoxy-2-tritertbutoxysiloxyethylene.

Example 7

The reaction was run as described in Example 5 with the exception that triethoxysilane was used as the starting silane. GC analysis showed that two major products had been formed approximately in a 1 to 1 ratio. GC/MS and NMR confirmed that the structures were 1,1-dimethyl-2-methoxy-1-triethoxysiloxyethylene and methyl 2-methyl-3-triethoxysilylpropanoate.

Example 8

The reaction was run as described in Example 5 with the exception that trimethoxysilane was used as the starting silane. GC analysis showed that two major products had been formed in approximately a 1 to 1 ratio. GC/MS and NMR confirmed that the structures were 1,1-dimethyl-2-methoxy-2-trimethoxysiloxyethylene and methyl 2-methyl-3-trimethoxysilylpropanoate.

Example 9

A 100 ml 3-neck flask equipped with a stirring bar, thermometer, 10 ml addition funnel,

and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 7.0g (0.028 mol) tritertbutoxysilane, 9.5g xylene, .40g phenothiazine and 200 ppm (10.37 mg) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to reflux at which time 2.5g (0.028 mol) methyl acrylate was added to the reaction mixture over a period of 1-2 hours which was then heated an additional 5 hours. The reaction was cooled to room temperature and GC analysis showed one major product had been formed in greater than 90% yield. GC/MS and NMR confirmed the structure to be 1-methyl-2-methoxy-2-tritertbutoxysiloxyethylene.

Example 10

The reaction was run as described in Example 9 with the exception that triisopropoxysilane was used as the starting silane. GC/MS and NMR confirmed the structure of the product formed to be 1-methyl-2-methoxy-2-triisopropoxysiloxyethylene.

Example 11

The reaction was run as described in Example 9 with the exception that triethoxysilane was used as the starting silane. GC analysis showed that two major products had been formed. GC/MS and NMR confirmed the structures to be 1-methyl-2-methoxy-2-triethoxysiloxyethylene and methyl 3-triethoxysilylpropanoate.

Example 12

The reaction was run as described in Example 9 with the exception that trimethoxysilane

was used as the starting silane. GC analysis showed that two major products had been formed. GC/MS and NMR confirmed the structures to be 1-methyl-2-methoxy-2-trimethoxysiloxyethylene and methyl 3-trimethoxysilylpropanoate.

Comparative A

The reaction was run as described in Example 5 except that tritertbutoxysilane was used as the starting silane and H_2PtCl_6 (200 ppm) was used as the catalyst. After heating the reaction mixture for 8 hours at 150°C no products were observed by GC.

Comparative B

The reaction was run as described in Example 13 except that H_2PtCl_6 (100 ppm) was used as the catalyst. After heating the reaction mixture for 8 hours at 150°C no products had been formed.

Example 13

A 50 ml 3-neck flask was equipped with a stirring bar, thermometer, 10 ml addition funnel and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 4.7g (0.029 mol) tris(dimethylamino)silane, 5.0g xylene, 0.28g phenothiazine and 200 ppm (8.32 mg) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to reflux, at which time 2.5g (0.029 mol) methyl acrylate was added dropwise to the reaction mixture over a period of 2 hours, which was then heated an additional 10 hours at 150°C. The reaction mixture was cooled to room temperature and GC analysis showed that one major product had been formed in greater than 80%

yield. GC/MS and NMR confirmed the structure to be 1-methyl-2-methoxy-2-tris(dimethylamino)siloxyethylene.

Example 14

A 1000 ml 3-neck flask was equipped with a stirring bar, thermometer, 250 ml addition funnel and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 200g (1.25 mol) tris(dimethylamino)silane, 307g xylene, 12.0g phenothiazine and 200 ppm (354 mg) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to reflux, at which time 106.8 (1.24 mol) methyl acrylate was added dropwise to the reaction over a period of 2-3 hours which was then heated for an additional 10 hours at 150°C. The reaction mixture was then cooled to room temperature and GC analysis showed the formation of one major product in greater than 80% yield. This was then distilled at 140°C/27 mbar (20 mm Hg): GC/MS and NMR confirmed the structure to be 1-methyl-2-methoxy-2-tris(dimethylamino)siloxyethylene.

Example 15

A 250 ml 3-neck flask was equipped with a stirring bar, thermometer, 50 ml addition funnel and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 30g (0.186 mole) tris(dimethylamino)silane, 50g xylene, 0.02g MEHQ and 200 ppm (44.4 mg) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to reflux at which time 18.6g (0.186 mol) methyl methacrylate was added dropwise to the reaction over a period of 1-2 hours, which was then heated an additional 8 hours at 150°C. The

reaction mixture was cooled and GC showed that a single product had been formed in greater than 70% yield. GC/MS and NMR confirmed the structure to be 1,1-dimethyl-2-methoxy-2-tris(dimethylamino)siloxyethylene.

Example 16

The reaction was run as described in Example 5 except that methyldiethoxysilane was used as the starting silane. GC analysis after the reaction was complete showed that two products had been formed in a approximately a 3 to 2 ratio. NMR and GC/MS confirmed that the structure of the major product was 1,1-dimethyl-2-methoxy-2-diethoxymethylsiloxyethylene and the minor product was methyl 2-methyl-3-diethoxymethylsilylpropanoate.

Example 17

The reaction was run as described in Example 5 except that isobutylmethacrylate was used instead of methyl methacrylate. One major product was formed and NMR and GC/MS confirmed the structure to be 1,1-dimethyl-2-isobutoxy-2-triisopropoxysiloxyethylene.

Example 18

The reaction was run as described in example 8 except that isobutylmethacrylate was used instead of methyl methacrylate. Two products were formed in a 2 to 1 ratio. The major one was determined by GC/MS and NMR to be 1,1-dimethyl-2-isobutoxy-2-trimethoxysiloxyethylene. The minor one was isobutyl 2-methyl-3-trimethoxysilylpropanoate.

Example 19

The reaction was run as described in example 1 except that 2-trimethylsiloxyethyl-methacrylate was used instead of methyl methacrylate. One major product was formed. GC/MS and NMR confirmed the structure to be 1,1-dimethyl-2-(2'-trimethylsiloxyethoxy)-2-bis(trimethylsilyloxy)-methylsiloxyethylene.

Example 20

The reaction was run as described in Example 15 except that dimethylaminodimethylsilane was used instead of tris(dimethylamino)silane. One major product was formed and NMR and GC/MS confirmed the structure to be 1, 1-dimethyl-2-methoxy-2-dimethylaminodimethylsiloxyethylene.

Example 21

The reaction was run as described in Example 15 except that methylbis(dimethylamino)silane was used instead of tris(dimethylamino)silane. One major product was formed and NMR and GC/MS confirmed the structure to be 1,1-dimethyl-2-methoxy-2-methyl bis(dimethylamino)siloxyethylene.

Example 22

The reaction was run as described in example 5 except that dimethylethoxysilane was used as the starting silane. Two products were formed in a ratio of 3 to 1. The structure of the major product was confirmed by GC/MS and NMR to be 1,1-

dimethyl-2-methoxy-2-dimethylethoxysiloxyethylene.

Example 23

A sample of 1,1-dimethyl-2-methoxy-2-triethoxysiloxyethylene prepared as described in Example 7 was used as an initiator for Group Transfer Polymerization. The procedure followed was identical to that described in the Journal of the American Chemical Society, Vol. 105, page 5706, with the only exception being that 1,1-dimethyl-2-methoxy-2-triethoxysiloxyethylene was used instead of 1-methoxy-1-(trimethylsiloxy)-2-methyl-1-propene. Group Transfer Polymerization did occur.

Example 24

A sample of 1,1-dimethyl-2-methoxy-2-bis(trimethylsilyloxy)methylsiloxyethylene prepared as described in Example 1 was used as an initiator for Group Transfer Polymerization. The procedure followed was identical to that described in the Journal of the American Chemical Society, Vol. 105, page 5706, with the only exception being that 1,1-dimethyl-2-methoxy-2-bis(trimethylsilyloxy)-methylsiloxyethylene was used instead of 1-methoxy-1-(trimethylsiloxy)-2-methyl-1-propene. Group Transfer Polymerization did occur.

Example 25

A sample of 1,1-dimethyl-2-methoxy-2-methyl bis(dimethylamino)siloxyethylene prepared as described in Example 21 was used as an initiator for Group Transfer Polymerization. The procedure followed was identical to that described in the

D-

Journal of the American Chemical Society, Vol. 105, pge 5706, with the only exception being that 1,1-dimethyl-2-methoxy-2-methylbis(dimethylamino) siloxyethylene was used instead of 1-methoxy-1-(trimethylsiloxy)-2-methyl-1-propene. Group Transfer Polymerization did occur.

Example 26

A sample of 1,1-dimethyl-2-methoxy-2-methyldiethoxysiloxy-ethylene prepared as described in Example 16 was used as an initiator for Group Transfer Polymerization. The procedure followed was identical to that described in the Journal of the American Chemical Society, Vol. 105, page 5706, with the only exception being that 1,1-dimethyl-2-methoxy-2-methyldiethoxy-siloxy-ethylene was used instead of 1-methoxy-1-(trimethylsiloxy)-2-methyl-1-propene. Group Transfer Polymerization did occur.

Example 27

A sample of 1,1-dimethyl-2-methoxy-2-dimethylaminodimethylsiloxyethylene prepared as described in Example 20 was used as an initiator for Group Transfer Polymerization. The procedure followed was identical to that described in the Journal of American Chemical Society, Vol. 105, page 5706, with the only exception being that 1,1-dimethyl-1-methoxy-2-dimethylaminodimethylsiloxy-ethylene was used instead of 1-methoxy-1-(trimethylsiloxy)-2-methyl-1-propene. Group Transfer Polymerization did occur.

Example 28

A 250 ml 3-neck flask was equipped with a stirring bar, thermometer, 50 ml addition funnel and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 48 g (0.714 mol) of methyldiethoxysilane, 25 g of triisopropylbenzene, 0.32 g monomethyl hydroquinone and 300 ppm (0.129 g) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated at 80°C at which time 25 g (0.357 mol) of methylvinylketone was added over a period of 1-2 hours. The reaction was exothermic and during the course of the addition of methylvinylketone the reaction temperature increased to 160°C. The reaction was cooled to room temperature and GC analysis showed that one major product had been formed. GC/MS confirmed that the major product was a mixture of the cis and trans isomers of 3-(methyldiethoxysiloxy)but-2-ene.

Example 29

The reaction was run as described in Example 28 with the exception that dimethylethoxysilane was used as the starting silane. GC analysis showed one major product had been formed, which GC/MS confirmed was a mixture of the cis and trans isomers of 3-(dimethylethoxysiloxy)but-2-ene.

Example 30

The reaction was run as described in Example 28 with the exception that triethoxysilane was used as the starting silane. GC analysis showed the formation of one major product which GC/MS

confirmed to be a mixture of cis and trans
3-(triethoxysiloxy)but-2-ene.

Example 31

A 500 ml 3-neck flask was equipped with a stirring bar, thermometer, 100 ml addition funnel and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 145 g (0.71 mol) tris(dimethylamino)silane, 115 g xylene, 0.33 g monomethyl hydroquinone and 300 ppm (0.31 g) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to 95°C, at which time 50 g (0.71 mol) methylvinylketone was added dropwise over a period of 1-2 hours. There was an initial exotherm which increased the temperature to 140°C. After the addition was complete the reaction was cooled to room temperature and GC analysis showed one major product had been formed. GC/MS confirmed that the major product was a mixture of cis and trans
3-(tris(dimethylamino)siloxy)but-2-ene.

Example 32

The reaction was run as described in Example 31 with the exception that methylbis(dimethylamino)silane was used as the starting silane and triisopropylbenzene was used as the solvent. GC/MS confirmed that the single product formed was a mixture of cis and trans
3-(methylbis(dimethylamino)siloxy)but-2-ene.

Example 33

A 500 ml 3-neck flask was equipped with a stirring bar, thermometer, 100 ml addition funnel

and reflux condenser topped with a nitrogen inlet tube. The flask was charged with 158 g (0.714 mol) bis(trimethylsiloxy)methylsilane, 150 g xylene, 0.42 g monomethyl hydroquinone and 300 ppm (0.43 g) $\text{RhCl}(\text{PPh}_3)_3$. The solution was heated to 80°C at which time 50 g (0.714 mol) methylvinylketone was added over a period of one to two hours. There was an initial exotherm and the reaction temperature increased to 140°C. After the addition was complete, the reaction was cooled to room temperature and GC analysis showed one major product had been formed in greater than 90% yield. GC/MS confirmed that this product was a mixture of cis and trans 3-(bis(trimethylsiloxy)methylsiloxy)but-2-ene.

Example 34

A sample of 1,1-dimethyl-2-methoxy-2-dimethylethoxy-siloxyethylene prepared as described in Example 22 was used as an initiator for Group Transfer Polymerization. The procedure followed was identical to that described in the Journal of the American Chemical Society, Vol. 105, page 5706, with the only exception being that 1,1-dimethyl-2-methoxy-2-dimethylethoxysiloxyethylene was used instead of 1-methoxy-1-(trimethylsiloxy)-2-methyl-1-propene. Group Transfer Polymerization did occur.

PATENTANWÄLTE
WUESTHOFF-v. PECHMANN-BEHRENS-GOETZ
EUROPEAN PATENT ATTORNEYS


Union Carbide Corp.
Our File: EP-59 902

0184692
DR.-ING. FRANZ WUESTHOFF
DR. PHIL. FREDA WUESTHOFF (1927-1956)
DIPL.-ING. GERHARD PULS (1952-1971)
DIPL.-CHEM. DR. E. FREIHERR VON PECHMANN
DR.-ING. DIETER BEHRENS
DIPL.-ING.; DIPL.-WIRTSCH.-ING. RUPERT GOETZ
D-8000 MÜNCHEN 90
SCHWEIGERSTRASSE 2
TELEFON: (089) 66 10 51
TELEGRAMM: PROTECIPATENT
TELEX: 5 24 070
TELEFAX: VIA (089) 2 71 60 63 (111)


CLAIMS

1. O-silylated ketene acetals or enol ethers consisting essentially of:


(a) O-aminosilylated ketene acetals and enol ethers of the general formula:



(b) O-siloxanesilylated ketene acetals and enol ethers of the general formulae:



(c) O-alkoxysilylated ketene acetals and enol ethers of the general formula:



wherein R, R¹ and R² are individually alkyl or aryl group containing 1 to 8 carbon atoms and R can be hydrogen; R³, R⁴, R⁵, R⁶, R⁷, and R¹⁰ are individually alkyl groups containing 1 to 8 carbon atoms; R⁸ is hydrogen or an alkyl or aryl groups having 1 to 8 carbon atoms; Z is selected

from hydrogen, an alkyl group, an aryl group, an alkoxy group and an aryloxy group containing from 1 to 15 carbon atoms and a is 0 or 1, b 0 or 1, y has a value from 1 to 500, z has a value from 0 to 499, and, the sum of y+z equals from 1 to 500 with the proviso that if z is 0, a + b must equal at least 1; x is 1, 2 or 3; and v is 0, 1 or 2.

2. The products of claim 1 wherein R, R¹ and R² are individually methyl, ethyl or phenyl groups and x has a value of 3.


3. The products of claim 1 wherein R, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, and R¹⁰ are methyl groups.

4. The products of claim 1 wherein v is 0.


5. The products of claims 1 to 4 being 1,1-dimethyl-2-methoxy-2-bis(trimethylsilyloxy)-methylsiloxylethylene, 1-methyl-2-methoxy-2-bis(trimethylsilyloxy)methylsiloxylethylene, 1,1-dimethyl-2-methoxy-2-heptamethylcyclo-tetrasiloxylethylene, 1,3-bis(1,1-dimethyl-2-methoxy-2-ethenoxy)-1,1,3,3-tetramethyldisiloxane, 1,1-dimethyl-2-methoxy-2-triisopropoxysiloxy-ethylene, 1,1-dimethyl-2-methoxy-2-tritertbutoxy-siloxethylene, 1,1-dimethyl-2-methoxy-1-tri-ethoxysiloxyethylene, 1-methyl-2-methoxy-2-tri-tertbutoxysiloxyethylene, 1-methyl-2-methoxy-2-triisopropoxysiloxyethylen, 1-methyl-2-methoxy-2-tris(dimethylamino)siloxyethylene of 1,1-dimethyl-2-methoxy-2-tris(dimethylamino)-siloxyethylene.

6. A process for preparing the products of claims 1 to 5 which comprises a 1,4-hydrosilation of a hydrosilation composition selected from


(a) aminosilanes of the formula:




(b) siloxanes of the formula:



(c) alkoxysilanes of the formula:



with an α,β -unsaturated carbonyl compound of the formula:



in the presence of a rhodium catalyst.

7. The process of claim 6 wherein the catalyst concentration is 0.00001 to 0.5 mole percent

based on the hydrosilation composition.

8. The process of claims 6 or 7 wherein the hydrosilation occurs at a temperature greater than 40° C, preferably between 70 and 150° C.

9. The process of claims 6 to 8 wherein a solvent is employed.

10. The use of the O-silylated ketene acetal or enol ether of claims 1 to 5 as the group transfer initiator in a group transfer polymerization reaction.

8184



European Patent
Office

EUROPEAN SEARCH REPORT

0184692

Application number

EP 85 11 4703

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	GB-A-2 130 229 (GENERAL ELECTRIC CO.) * Page 1 *	1,4,6-10	C 07 F 7/18 C 07 F 7/04 C 07 F 7/02 C 07 F 7/21
X	US-A-2 917 530 (D.L. BAILEY) * Whole document *	1,3,5-10	
X	GB-A-1 044 448 (RHONE-POULENC SA) * Pages 1,2; page 3, lines 66-89; examples VIII, IX *	1,6-10	
A	US-A-3 450 737 (Y. COLLEVILLE) * Claims; examples *	1-10	
A	FR-A-2 187 798 (SAGAMI CHEMICAL RESEARCH CENTER) * Examples *	1-10	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
D,A	CHEMICAL ABSTRACTS, vol. 82, no. 15, 14th April 1975, page 444, no. 98038j, Columbus, Ohio, US; E. YOSHII et al.: "Hydrosilylation of alpha-beta-unsaturated esters", & CHEM. PHARM. BULL. 1974, 22(11) 2767-9 * Whole document *	2-10	C 07 F 7/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 03-03-1986	Examiner PAUWELS G.R.A.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO Form 1500 03 82



European Patent
Office

EUROPEAN SEARCH REPORT

0184692

Application number

EP 85 11 4703

DOCUMENTS CONSIDERED TO BE RELEVANT				Page 2																																			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)																																				
D, A	US-A-4 417 034 (O.W. WEBSTER) * Examples *	1-10																																					
D, A	--- JOURNAL OF ORGANOMETALLIC CHEMISTRY, vol. 111, 1976, pages 43-60, Elsevier Sequoia SA, NL; I. OJIMA et al.: "Hydrosilylation of alpha,beta-unsaturated nitriles and esters catalyzed by tris(triphenylphosphine)chlororho- dium" * Pages 47-49, 55-60 * -----	1-10																																					
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)																																				
The present search report has been drawn up for all claims																																							
Place of search THE HAGUE		Date of completion of the search 03-03-1986		Examiner PAUWELS G.R.A.																																			
<table border="0"><tr><td colspan="2">CATEGORY OF CITED DOCUMENTS</td><td colspan="3">T theory or principle underlying the invention</td></tr><tr><td>A</td><td>particularly relevant if taken alone</td><td>E</td><td colspan="2">earlier patent document, but published on, or</td></tr><tr><td>Y</td><td>particularly relevant if combined with another</td><td>D</td><td colspan="2">after the filing date</td></tr><tr><td></td><td>document of the same category</td><td>C</td><td colspan="2">document cited in the application</td></tr><tr><td>A</td><td>technological background</td><td>L</td><td colspan="2">document cited for other reasons</td></tr><tr><td>O</td><td>non-written disclosure</td><td>R</td><td colspan="2">member of the same patent family, corresponding</td></tr><tr><td>P</td><td>intermediate document</td><td></td><td colspan="2">document</td></tr></table>					CATEGORY OF CITED DOCUMENTS		T theory or principle underlying the invention			A	particularly relevant if taken alone	E	earlier patent document, but published on, or		Y	particularly relevant if combined with another	D	after the filing date			document of the same category	C	document cited in the application		A	technological background	L	document cited for other reasons		O	non-written disclosure	R	member of the same patent family, corresponding		P	intermediate document		document	
CATEGORY OF CITED DOCUMENTS		T theory or principle underlying the invention																																					
A	particularly relevant if taken alone	E	earlier patent document, but published on, or																																				
Y	particularly relevant if combined with another	D	after the filing date																																				
	document of the same category	C	document cited in the application																																				
A	technological background	L	document cited for other reasons																																				
O	non-written disclosure	R	member of the same patent family, corresponding																																				
P	intermediate document		document																																				

419,6 Jun. 1987 03 82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.